

Predicting positivity for a new era of Alzheimer disease prevention trials

Jeffrey M. Burns, MD,
MS
William E. Klunk, MD,
PhD

Correspondence & reprint
requests to Dr. Burns:
jburns2@kumc.edu

Neurology® 2012;79:1530–1531

Alzheimer disease (AD) pathophysiology likely begins years prior to the emergence of clinical symptoms. Biomarker studies suggest deposition of cerebral amyloid may be a necessary and early component of the AD pathophysiologic process.¹ The advent of amyloid imaging techniques with the introduction of Pittsburgh compound B (PiB) in 2004² promised the identification of individuals in the presumptive presymptomatic stages of AD.

Autopsy studies have shown that about one-third of older adults have substantial AD neuropathologic changes without cognitive or functional manifestations. Accordingly, amyloid imaging studies demonstrated the presence of cerebral amyloid deposits in 20% to 40% of cognitively normal older adults, although at levels typically less than those observed in individuals with AD. Research criteria³ were introduced in 2011, defining cognitively normal individuals who test positive for the presence of cerebral amyloid as having “preclinical AD,” and plans are underway to target these individuals in large-scale trials assessing therapeutic interventions and their modification of dementia risk.

It is with these future trials in mind that Mielke et al.,⁴ in this issue of *Neurology*®, examined common and noninvasive screening measures that might help identify individuals with elevated cerebral amyloid. Their goal was to inform clinical trial design by developing a relatively inexpensive method to enrich a sample of cognitively normal individuals for amyloid positivity and thus reduce the number of individuals who need to be screened. This is an important goal as the coming wave of prevention trials will need to screen out up to 80% of potential participants who are amyloid negative, at a considerable expense and burden that could limit the size and power of such studies. To put this in perspective, in a trial such as the Anti-Amyloid Treatment in Asymptomatic Alzheimer Disease (A4) using amyloid-PET to screen 3,200 amyloid-negative subjects so as to arrive at ~800 amyloid-positive subjects (400 each in placebo

and treatment arms), screening would add approximately \$8,000,000 to the cost of the trial.

Mielke et al. examined a large number ($n = 483$) of cognitively normal older adults (ages 70–92 years) from a population-based cohort characterized by PiB-PET. Participants were classified as being amyloid-positive using 2 different cutpoints, with 44% of the group classified as amyloid-positive using the more lenient cutpoint (cortical to cerebellar PiB uptake >1.4) and 31% amyloid-positive using a higher cutpoint (>1.5). The investigators tested the predictive value of age, sex, *APOE* genotype, family history, cognitive performance, and subjective cognitive complaints.

Not surprisingly, given findings from prior studies,⁵ the best indicators of the presence of cerebral amyloid were age and *APOE* $\epsilon 4$. For every 5-year increase in age, the odds of being amyloid-positive increased 40%–45%, while *APOE* $\epsilon 4$ carriers were 3 times more likely to be amyloid-positive. Other significant predictors of cerebral amyloid positivity included family history of AD, reduced cognitive performance on an extensive neuropsychological battery, and subjective memory complaints. Interestingly, subjective memory complaints were as good at predicting amyloid presence as the extensive battery of cognitive tests, suggesting that assessing memory complaints is as effective as cognitive performance for predicting amyloid positivity. Additionally, the data suggest that the predictive value of the variables (with the exception of cognitive performance) appears to be attenuated in the older age group (>80 years), further underscoring the importance of age in screening for amyloid positivity.

The main limitation to the practical value of these findings is that using the best-performing predictors to enrich clinical trial samples—advanced age and *APOE* $\epsilon 4$ carrier status—would severely limit the generalizability of the findings. For instance, limiting trials to subjects over age 80 or to $\epsilon 4$ -positive subjects would miss a sizable target population and, as

See page 1570

From the Department of Neurology (J.M.B.), University of Kansas Alzheimer's Disease Center, University of Kansas School of Medicine, Kansas City; and Departments of Psychiatry and Neurology (W.E.K.), University of Pittsburgh School of Medicine, Pittsburgh, PA.
Go to Neurology.org for full disclosures. Disclosures deemed relevant by the authors, if any, are provided at the end of this editorial.

the authors note, restrict the labeling uses of any new therapy granted by a regulatory agency. An additional limitation is the exclusion of participants under the age of 70 years; it is in this age range that screening methods may have the biggest effect on enriching trial populations, given the lower prevalence of amyloidosis.

US Food and Drug Administration approval of the ^{18}F -based amyloid imaging agent florbetapir this year will ensure widespread availability of amyloid imaging and could further stimulate a wave of AD prevention trials. Nonetheless, it is important to note that the clinical importance of asymptomatic amyloid plaques remains imprecisely defined. Although the concept of preclinical AD posits that cerebral amyloid deposition in cognitively normal adults represents a presymptomatic stage of AD, autopsy studies clearly demonstrate that not all amyloid-positive individuals will develop a dementia syndrome prior to their death. Additionally, the magnitude and timing of risk associated with asymptomatic cerebral amyloidosis is not yet well-defined. Without clear knowledge of these risks, and in the absence of effective interventions to delay the onset of dementia, the use of amyloid imaging in unaffected, cognitively normal participants should be restricted to the research arena, such as is planned for the prevention trials. Additionally, since it is unlikely that amyloid-negative subjects will be included in prevention trials, participation in these trials will reveal the participant's positive amyloid status. This requires careful consideration of how participants will be informed and counseled about this information.

Although there remain a number of important unknowns, early studies suggest the presence of cerebral amyloid is not benign and thus support the use of amyloid imaging for identifying an important target cohort for AD prevention trials. For instance, results from early studies suggest that asymptomatic cerebral amyloid is associated with cognitive decline,⁶ brain atrophy,⁷ and altered brain function,^{8,9} supporting the concept of targeting this population for testing dementia-delaying strategies. The findings from the Mielke et al. study suggest that inexpensive and noninvasive measures can be used to reduce the number of cognitively normal individuals who need

to be screened for these trials; the data thereby represent useful information for ushering in a new era of AD prevention studies.

DISCLOSURE

J.M. Burns receives research support from the NIH; has served on the Novartis speakers' bureau and as a consultant for Johnson County Clinical Trials and PRA International; receives research support from Jansen Corporation, Wyeth, Danone, and Baxter; and has served as an expert witness in legal proceedings regarding competency. W.E. Klunk has received support from NIH and has received consulting fees from GE Healthcare, Janssen AI, Elan, Hoffmann-LaRoche, Wyeth/Pfizer, and Eli Lilly. GE Healthcare holds a license agreement with the University of Pittsburgh based on the technology described in this manuscript. Dr. Klunk is a coinventor of PiB and, as such, has a financial interest in this license agreement. GE Healthcare provided no grant support for this study and had no role in the design or interpretation of results or preparation of this manuscript. **Go to Neurology.org for full disclosures.**

REFERENCES

1. Jack CR Jr, Knopman DS, Jagust WJ, et al. Hypothetical model of dynamic biomarkers of the Alzheimer's pathological cascade. *Lancet Neurol* 2010;9:119–128.
2. Klunk WE, Engler H, Nordberg A, et al. Imaging brain amyloid in Alzheimer's disease with Pittsburgh compound-B. *Ann Neurol* 2004;55:306–319.
3. Sperling RA, Aisen PS, Beckett LA, et al. Toward defining the preclinical stages of Alzheimer's disease: recommendations from the National Institute on Aging–Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimers Dement* 2011;7:280–292.
4. Mielke MM, Wiste HJ, Weigand SD, et al. Indicators of amyloid burden in a population-based study of cognitively normal elderly. *Neurology* 2012;79:1570–1577.
5. Morris JC, Roe CM, Xiong C, et al. APOE predicts amyloid-beta but not tau Alzheimer pathology in cognitively normal aging. *Ann Neurol* 2010;67:122–131.
6. Resnick SM, Sojkova J, Zhou Y, et al. Longitudinal cognitive decline is associated with fibrillar amyloid-beta measured by [^{11}C]PiB. *Neurology* 2010;74:807–815.
7. Storandt M, Mintun MA, Head D, Morris JC. Cognitive decline and brain volume loss as signatures of cerebral amyloid-beta peptide deposition identified with Pittsburgh compound B: cognitive decline associated with Abeta deposition. *Arch Neurol* 2009;66:1476–1481.
8. Sperling RA, Laviolette PS, O'Keefe K, et al. Amyloid deposition is associated with impaired default network function in older persons without dementia. *Neuron* 2009;63:178–188.
9. Hedden T, Van Dijk KR, Becker JA, et al. Disruption of functional connectivity in clinically normal older adults harboring amyloid burden. *J Neurosci* 2009;29:12686–12694.